

IN THE CLAIMS:

Please cancel Claims 2, 8, 17, 18 and 20.

1. (Currently Amended) A process for copolymerizing polar and non-polar monomers, characterized in that at least one polar monomer selected from the group consisting of chloroprene, styrene, acrylonitrile, vinyl chloride, acrylic acid, acrylate, cyanacrylate, methacrylic acid, methacrylate, acrylamide, methacrylonitrile, vinyl acetate, propene oxide, ethene oxide, vinyl carbazole, vinylpyrrolidone, vinyl ester, and compounds built up therefrom and at least one non-polar monomer selected from the group consisting of olefin, diolefin and triene are polymerized in the presence of one or more transition metal compounds conforming structurally to ML_aQ_b wherein M is a metal selected from the group consisting of chromium, manganese, iron, cobalt, nickel, ruthenium, rhodium and palladium, L is a 2-, 3- or 4-dentate chelating ligand, Q is a mono-anionic or non-ionic ligand, b is an integer equal to or greater than 1 and a is a positive number calculated as the total number of receptor coordination sites on M - b / the number of donor coordination sites on the ligand, one or more radical-producers and optionally one or more co-catalysts[.];
and
the transition metal compound is chosen so that the transition metal compound, optionally in the presence of a co-catalyst, reversibly forms a complex with the radically growing polymer chain and the said non-polar monomers are inserted into the bond thus formed between transition metal and polymer chain.

2. (Cancelled)

3. (Previously Presented) A process according to Claim 1 characterized in that the radical-producer is chosen so that the radical-producer(s) initiate polymerization and do not react in a detrimental fashion with the transition metal compound.

4. (Currently Amended) ~~A process according to Claim 1 characterized in that~~ A process for copolymerizing polar and non-polar monomers, characterized in that at least one polar monomer selected from the group consisting of chloroprene,

Mo-6856

-2-

styrene, acrylonitrile, vinyl chloride, acrylic acid, acrylate, cyanacrylate, methacrylic acid, methacrylate, acrylamide, methacrylonitrile, vinyl acetate, propene oxide, ethene oxide, vinyl carbazole, vinylpyrrolidone, vinyl ester, and compounds built up therefrom and at least one non-polar monomer selected from the group consisting of olefin, diolefin and triene are polymerized in the presence of one or more transition metal compounds conforming structurally to ML_aQ_b wherein M is a metal selected from the group consisting of chromium, manganese, iron, cobalt, nickel, ruthenium, rhodium and palladium, L is a 2-, 3- or 4-dentate chelating ligand, Q is a mono-anionic or non-ionic ligand, b is an integer equal to or greater than 1 and a is a positive number calculated as the total number of receptor coordination sites on M - b) / the number of donor coordination sites on the ligand, one or more radical-producers and optionally one or more co-catalysts; and
one or more transition metal complex cation forming compounds or coordination complex compounds are used as co-catalyst, chosen from the group of strong, neutral Lewis acids, ionic compounds with Lewis acid cations or Broenstedt acid cations and non-coordinating anions.

5. (Currently Amended) A composition containing one or more transition metal compounds conforming structurally to ML_aQ_b wherein M is a metal selected from the group consisting of chromium, manganese, iron, cobalt, nickel, ruthenium, rhodium and palladium, L is a 2-, 3- or 4-dentate chelating ligand, Q is a mono-anionic or non-ionic ligand, b is an integer equal to or greater than 1 and a is a positive number calculated as the total number of receptor coordination sites on M - b) / the number of donor coordination sites on the ligand, one or more radical-producers and optionally one or more co-catalyst[[s.]]; and one or more compounds chosen from the group of strong, neutral Lewis acids, ionic compounds with Lewis acid cations or Broenstedt acid cations and non-coordinating anions are used as co-catalysts.

6. (Cancelled)

7. (Previously Presented) A composition according to Claim 5 characterized in that the radical-producer is a peroxide, a diazo compound or a mixture thereof.

8. (Cancelled)

9. (Currently Amended) ~~A composition according to Claim 5~~
characterized in that A composition containing one or more transition metal compounds conforming structurally to ML_aQ_b , wherein M is a metal selected from the group consisting of chromium, manganese, iron, cobalt, nickel, ruthenium, rhodium and palladium, L is a 2-, 3- or 4-dentate chelating ligand, Q is a mono-anionic or non-ionic ligand, b is an integer equal to or greater than 1 and a is a positive number calculated as the total number of receptor coordination sites on M - b) / the number of donor coordination sites on the ligand, one or more radical-producers and optionally one or more co-catalyst; and
wherein the transition metal compound is chosen so that the transition metal compound, optionally in the presence of a co-catalyst, can reversibly form a complex with a radically growing polymer chain and non-polar monomers selected from the group consisting of olefin, diolefin and triene can be inserted into the bond formed in this way between transition metal and polymer chain.

10. (Previously Presented) A method of using the composition according to Claim 5 comprising catalyzing the copolymerization of monomers.

11. (Previously Amended) Copolymers which have a statistical distribution on the molecular level prepared in a process according to Claim 1.

12. (Cancelled)

13. (Previously Presented) A process for producing a copolymer comprising polymerizing at least one polar monomer selected from the group consisting of chloroprene, styrene, acrylonitrile, vinyl chloride, acrylic acid, acrylate, cyanacrylate, methacrylic acid, methacrylate, acrylamide, methacrylonitrile, vinyl

Mo-6856

acetate, propene oxide, ethene oxide, vinyl carbazole, vinylpyrrolidone, vinyl ester, and compounds built up therefrom and at least one non-polar monomer selected from the group consisting of olefin, diolefin and triene in the presence of one or more compounds of transition metals conforming structurally to ML_aQ_b wherein M is a metal selected from the group consisting of chromium, manganese, iron, cobalt, nickel, ruthenium, rhodium and palladium, L is a 2-, 3- or 4-dentate chelating ligand, Q is a mono-anionic or non-ionic ligand, b is an integer equal to or greater than 1 and a is a positive number calculated as the total number of receptor coordination sites on M - b) / the number of donor coordination sites on the ligand and one or more radical-producers.

14. (Previously Presented) The process of Claim 13 wherein the polymerizing is in the further presence of one or more co-catalysts.

15. (Previously Presented) The copolymer prepared by the process of Claim 13.

16. (Currently Amended) A composition containing one or more transition metal compounds conforming structurally to ML_aQ_b wherein M is a metal selected from the group consisting of chromium, manganese, iron, cobalt, nickel, ruthenium, rhodium and palladium, L is a 2-, 3- or 4-dentate chelating ligand, Q is a mono-anionic or non-ionic ligand, b is an integer equal to or greater than 1 and a is a positive number calculated as the total number of receptor coordination sites on M - b) / the number of donor coordination sites on the ligand and one or more radical-producers[.]; and
at least one co-catalyst selected from the group consisting of strong Lewis acids, neutral Lewis acids, ionic compounds with Lewis acid cations, ionic compounds with Brønstedt acid cations, and non-coordinating anions.

17. (Cancelled)

18. (Cancelled)

Mo-6856

-5-

19. (Previously Presented) The composition of Claim 16 wherein the radical-producer is at least one member selected from the group consisting of peroxide and a diazo compound.

20. (Cancelled)

21. (Previously Presented) A method of using the copolymer of Claim 15 comprising preparing a molded article.

22. (Cancelled)

23. (Previously Presented) A method of using a copolymer the production of which comprises polymerizing at least one polar monomer selected from the group consisting of chloroprene, styrene, acrylonitrile, vinyl chloride, acrylic acid, acrylate, cyanacrylate, methacrylic acid, methacrylate, acrylamide, methacrylonitrile, vinyl acetate, propene oxide, ethene oxide, vinyl carbazole, vinylpyrrolidone, vinyl ester, and compounds built up therefrom and at least one non-polar monomer selected from the group consisting of olefin, diolefin and triene in the presence of one or more compounds of transition metals conforming structurally to ML_aQ_b wherein M is a metal selected from the group consisting of chromium, manganese, iron, cobalt, nickel, ruthenium, rhodium and palladium, L is a 2-, 3- or 4-dentate chelating ligand, Q is a mono-anionic or non-ionic ligand, b is an integer equal to or greater than 1 and a is a positive number calculated as the total number of receptor coordination sites on M - b) / the number of donor coordination sites on the ligand and one or more radical-producers, the method comprising preparing an adhesive.